## Lactation Management Self-Study Modules, Level I Post-Test (without answers)

Please circle the appropriate response:

1.	Identify the componer	nt of human milk	that binds iron I	ocally to in	ihibit bacterial	growth:
	racinary and compensor	it of floridge from		Occurry to it	II IIDIK DUQUURI	4101

- a. taurine
- b. secretory IgA
- c. macrophages
- d. lactoferrin
- e. oligosaccharides
- 2. Identify the component of human milk that provides specific immunity against many organisms:
  - a. taurine
  - b. secretory IgA
  - c. macrophages
  - d. lactoferrin
  - e. oligosaccharides
- 3. The <u>most important criterion</u> for assessing the milk transfer during a feeding at the breast is:
  - a. visible areola compression
  - b. audible swallow
  - c. proper alignment
  - d. proper attachment
- 4. Compared to formula, human milk contains higher levels of:
  - a. vitamin D
  - b. iron
  - c. lipase
  - d. vitamin A
  - e. none of the above
- 5. The hormone considered responsible for milk ejection is:
  - a. progesterone
  - b. prolactin
  - c. estrogen
  - d. oxytocin

- 6. A mother with a three-day old baby presents with sore nipples. The problem began with the first feeding and has persisted with every feeding. The most likely source of the problem is:
  - a. feeding too long
  - b. poor attachment
  - c. baby's suck is too strong
  - d. lack of nipple preparation during pregnancy
- 7. The hormone considered responsible for milk synthesis is:
  - a. progesterone
  - b. prolactin
  - c. estrogen
  - d. oxytocin
- 8. Which of the following would you suggest that a woman with inverted nipples do during the third trimester?
  - a. Use breast shells with guidance from her health care provider
  - b. Cut holes in the bra to allow the nipples to protrude; wear it day and night
  - c. Encourage everting the nipples four times a day to permanently evert her nipples
  - d. Do nothing because the natural changes in the breast during pregnancy and the infant's suckling postpartum may evert the nipples
- 9. Which of the following is most likely to have the greatest effect on the volume of milk a woman produces?
  - a. maternal weight for height
  - b. maternal fluid intake
  - c. supplementation of the infant with formula
  - d. maternal caloric intake
  - e. both a and c
- 10. Infants exclusively breastfed for about six months will have:
  - a. Fewer episodes of lower respiratory infection
  - b. fewer episodes of diarrhea
  - c. none of the above
  - d. both a and b above
- 11. The addition of complementary foods to breastfed infants is recommended at about:
  - a. 2 months
  - b. 4 months
  - c. 6 months
  - d. 8 months
  - e. 10 months

- 12. Signs of adequate breast milk intake in the early (first 4-6) weeks include all EXCEPT:
  - a. baby gains weight
  - b. at least 3-4 stools in 24 hours
  - c. sounds of swallowing
  - d. baby sleeps through the night
  - e. at least 6 diapers wet with urine in 24 hours
- 13. It is especially important that an infant with a strong family history of allergy should be exclusively breastfed for:
  - a. 2 months
  - b. 4 months
  - c. 6 months
  - d. 8 months
  - e. 10 months
- 14. Severe engorgement is most often due to:
  - a. high oxytocin level
  - b. infrequent feedings
  - c. high prolactin level
  - d. postpartum depression
- 15. The most common cause of poor weight gain among breastfed infants during the first four weeks after birth is:
  - a. maternal endocrine problems
  - b. maternal nutritional deficiencies
  - c. infant metabolic disorders
  - d. infrequent or ineffective feedings
  - e. low fat content of breast milk
- 16. A breastfeeding mother with a 3-month old infant has a red tender wedge-shaped area on the outer quadrant of one breast. She has flu-like symptoms and a temperature of 39° C. Your management includes all of the following EXCEPT:
  - a. extra rest
  - b. interrupt breastfeeding for 48 hours
  - c. moist heat to the involved region
  - d. antibiotics for 10 to 14 days days
- 17. Studies have indicated that the Lactational Amenorrhea Method (LAM) of contraception is less reliable under which of the following circumstances:
  - a. feeds 8 or more times in 24 hours
  - b. is given no regular supplements
  - c. is less than 8 months old
  - d. continues with night feedings

- 18. Which of the following statements is not true of The International Code of Marketing of Breastmilk Substitutes approved as a resolution in the World Health Assembly (WHA) in 1981:
  - a. is updated every two years by the WHA
  - b. provides guidelines for the ethical marketing of infant formula
  - c. is incorporated into the Baby Friendly Hospital assessment
  - d. was approved by all WHA member countries
  - e. includes bottles, nipples, and breastmilk substitutes
- 19. Nipple candidiasis can be associated with all of the following EXCEPT:
  - a. oral thrush in the infant
  - b. burning pain in the breast
  - c. fever and malaise
  - d. pink and shiny appearance of the nipples and areola
- 20. Jaundice in a normal full term breastfeeding infant is improved by:
  - a. giving glucose water after breastfeeding
  - b. giving water after breastfeeding
  - c. breastfeeding frequently (at least 8 or more times in 24 hours)
  - d. both a and c
- 21. Breastfeeding is contraindicated in which of the following conditions:
  - a. infant with galactosemia
  - b. mother with mastitis
  - c. mother with hepatitis B
  - d. mother with inverted nipples
  - e. both a and c
- 22. Reasons for including breastfeeding support for mother infant in planning for or responding to major emergencies where clean water, sanitation and power are disrupted do not include:
  - a. It is less expensive than providing for infant formula
  - b. With support even mothers who have already weaned can be assisted to relactate
  - c. Breastmilk provides immunoglobulins that actively prevent infection.
  - d. In a stressful emergency situation breastfeeding provides a secure environment for infants and young children
- 23. Hospital policies that promote breastfeeding include:
  - a. use of a dropper for routine water supplementation
  - b. uninterrupted sleep the first night to allow mother's milk supply to build up
  - c. unlimited access of mother to baby
  - d. use of pacifiers to prevent sore nipples

24. through 28. Label the structures of the breast by inserting next to the appropriate pointer the number of the structure listed below:

24. Montgomery's glands25. Supporting fat and other tissues26. Alveoli

27. Areola

28. Duct

